

Z(3)-symmetric effective theory for pure gauge QCD at high temperature

Alexi Vuorinen

University of Washington, Seattle

PRD 74: 025011 (2006), hep-ph/0604100 with Larry Yaffe

Outline

- Background
 - Dimensional reduction and QCD
 - The center symmetry and the Wilson line
- Construction of the new theory
 - Degrees of freedom, potentials
 - Perturbative matching to full theory
 - The $Z(3)$ domain wall
- Phase diagram of the new theory
- Conclusions and future directions

QCD and dimensional reduction

- Conventional DR: at high $T \gg gT$, integrate out all non-static modes ($m \sim 2\pi T$) to obtain 3d effective theory for the static modes

$$\begin{aligned}\mathcal{L}_{\text{EQCD}} = & g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} [(D_i A_0)^2] \right. \\ & \left. + m_E^2 \text{Tr} (A_0^2) + \lambda_E \text{Tr} (A_0^4) \right\} + \delta \mathcal{L}_E, \\ & g_3 \equiv \sqrt{T} g, \quad m_E \sim gT, \quad \lambda_E \sim g^2\end{aligned}$$

QCD and dimensional reduction

- Conventional DR: at high $T \gg gT$, integrate out all non-static modes ($m \sim 2\pi T$) to obtain 3d effective theory for the static modes

$$\mathcal{L}_{\text{EQCD}} = g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} [(D_i A_0)^2] \right. \\ \left. + m_E^2 \text{Tr} (A_0^2) + \lambda_E \text{Tr} (A_0^4) \right\} + \delta \mathcal{L}_E,$$

$$g_3 \equiv \sqrt{T} g, \quad m_E \sim gT, \quad \lambda_E \sim g^2$$

- New theory sufficient to describe equilibrium thermodynamics at length scales $\gtrsim 1/(gT)$

QCD and dimensional reduction


- Conventional DR: at high $T \gg gT$, integrate out all non-static modes ($m \sim 2\pi T$) to obtain 3d effective theory for the static modes

$$\mathcal{L}_{\text{EQCD}} = g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} [(D_i A_0)^2] \right. \\ \left. + m_E^2 \text{Tr} (A_0^2) + \lambda_E \text{Tr} (A_0^4) \right\} + \delta \mathcal{L}_E,$$


$$g_3 \equiv \sqrt{T} g, \quad m_E \sim gT, \quad \lambda_E \sim g^2$$

- New theory sufficient to describe equilibrium thermodynamics at length scales $\gtrsim 1/(gT)$
- Parameters available through comparison of long distance correlators in EQCD and full QCD


- Partial success: impressive perturbative results derived at high T



- Partial success: impressive perturbative results derived at high T
 - Problems in non-perturbative regime



- Partial success: impressive perturbative results derived at high T
 - Problems in non-perturbative regime



- Fundamental problem: all symmetries of original theory are *not* respected by the reduction!

The center symmetry

- Full gauge symmetry of SU(3) Yang-Mills theory

$$A_\mu(x) \rightarrow s(x) (A_\mu(x) + i \partial_\mu) s(x)^\dagger, s(x) \in SU(3)$$
$$s(x + \beta \hat{e}_t) = z s(x), z \in Z(3)$$

The center symmetry

- Full gauge symmetry of SU(3) Yang-Mills theory

$$A_\mu(x) \rightarrow s(x) (A_\mu(x) + i \partial_\mu) s(x)^\dagger, s(x) \in SU(3)$$
$$s(x + \beta \hat{e}_t) = z s(x), z \in Z(3)$$

under which the Wilson line transforms as a Z(3) fundamental

$$\Omega(\mathbf{x}) \equiv \mathcal{P} \exp \left[i \int_0^\beta d\tau A_0(\tau, \mathbf{x}) \right]$$
$$\text{Tr } \Omega(\mathbf{x}) \rightarrow z \text{Tr } \Omega(\mathbf{x})$$

The center symmetry

- Full gauge symmetry of SU(3) Yang-Mills theory

$$A_\mu(x) \rightarrow s(x) (A_\mu(x) + i \partial_\mu) s(x)^\dagger, s(x) \in SU(3)$$
$$s(x + \beta \hat{e}_t) = z s(x), z \in Z(3)$$

under which the Wilson line transforms as a Z(3) fundamental

$$\Omega(\mathbf{x}) \equiv \mathcal{P} \exp \left[i \int_0^\beta d\tau A_0(\tau, \mathbf{x}) \right]$$

$$\text{Tr } \Omega(\mathbf{x}) \rightarrow z \text{Tr } \Omega(\mathbf{x})$$

- Ω order parameter for deconfinement transition
 - $|\langle \text{Tr } \Omega(\mathbf{x}) \rangle| = e^{-\beta \Delta F_q(\mathbf{x})}$

- In deconfined phase, effective potential for Ω has degenerate minima $\Omega_{\min} \sim e^{i2\pi n/3} \mathbf{1}$, $n \in \{0, 1, 2\}$
 - Tunnelings between different vacua important near T_c
 - At (1st order) phase transition quadruple point with phase coexistence with the confining one

- In deconfined phase, effective potential for Ω has degenerate minima $\Omega_{\min} \sim e^{i2\pi n/3} \mathbf{1}$, $n \in \{0, 1, 2\}$
 - Tunnelings between different vacua important near T_c
 - At (1st order) phase transition quadruple point with phase coexistence with the confining one
- EQCD Lagrangian derived expanding effective potential around $A_0 = 0$
 - $Z(3)$ invariance lost
 - Complex $Z(3)$ minima $A_0 = \frac{2\pi T}{3}$ completely outside the domain of validity of eff. theory
 - (One important) cause of problems in EQCD phase diagram and predictions near T_c

$Z(3)$ invariant theory

- Want to build a superrenormalizable $3d$ effective theory that
 - Reduces to EQCD at high T
 - Respects $Z(3)$, correct domain wall physics

Z(3) invariant theory

- Want to build a superrenormalizable 3d effective theory that
 - Reduces to EQCD at high T
 - Respects Z(3), correct domain wall physics
- Minimal set of dof's: A_i and Ω
 - $\Omega \in \text{SU}(3) \Rightarrow$ polynomial Lagrangian non-renormalizable

Z(3) invariant theory

- Want to build a superrenormalizable 3d effective theory that
 - Reduces to EQCD at high T
 - Respects Z(3), correct domain wall physics
- Minimal set of dof's: A_i and Ω
 - $\Omega \in \text{SU}(3) \Rightarrow$ polynomial Lagrangian non-renormalizable

Sigma Models	
Non-linear	Linear
$\bar{\phi} \cdot \phi = \mathbb{1}$	Polynomial V
Same long distance physics!	

Z(3) invariant theory

- Want to build a superrenormalizable $3d$ effective theory that
 - Reduces to EQCD at high T
 - Respects Z(3), correct domain wall physics
- Minimal set of dof's: A_i and Ω
 - $\Omega \in \text{SU}(3) \Rightarrow$ polynomial Lagrangian non-renormalizable
- New (old) idea: replace Ω by $\mathcal{Z} \in \text{GL}(3, \mathbb{C})$
 - Coarse-grained version of Ω
 - After gauge fixing, contains $10 - 2 = 8$ unphysical dof's that are chosen heavier ($m \sim T$) than the physical ones ($m \lesssim gT$)

- Require gauge and $Z(3)$ invariance

$$\mathcal{Z}(\mathbf{x}) \rightarrow s(\mathbf{x}) \mathcal{Z}(\mathbf{x}) s(\mathbf{x})^\dagger,$$

$$\mathbf{A}(\mathbf{x}) \rightarrow s(\mathbf{x}) (\mathbf{A}(\mathbf{x}) + i\nabla) s(\mathbf{x})^\dagger,$$

$$\mathcal{Z}(\mathbf{x}) \rightarrow e^{2\pi i n/3} \mathcal{Z}(\mathbf{x})$$

- Require gauge and $Z(3)$ invariance

$$\mathcal{Z}(\mathbf{x}) \rightarrow s(\mathbf{x}) \mathcal{Z}(\mathbf{x}) s(\mathbf{x})^\dagger,$$

$$\mathbf{A}(\mathbf{x}) \rightarrow s(\mathbf{x}) (\mathbf{A}(\mathbf{x}) + i\nabla) s(\mathbf{x})^\dagger,$$

$$\mathcal{Z}(\mathbf{x}) \rightarrow e^{2\pi i n/3} \mathcal{Z}(\mathbf{x})$$

and compose Lagrangian as

$$\mathcal{L} = g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} (D_i \mathcal{Z}^\dagger D_i \mathcal{Z}) + V(\mathcal{Z}) \right\},$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

- Require gauge and $Z(3)$ invariance

$$\mathcal{Z}(\mathbf{x}) \rightarrow s(\mathbf{x}) \mathcal{Z}(\mathbf{x}) s(\mathbf{x})^\dagger,$$

$$\mathbf{A}(\mathbf{x}) \rightarrow s(\mathbf{x}) (\mathbf{A}(\mathbf{x}) + i\nabla) s(\mathbf{x})^\dagger,$$

$$\mathcal{Z}(\mathbf{x}) \rightarrow e^{2\pi i n/3} \mathcal{Z}(\mathbf{x})$$

and compose Lagrangian as

$$\mathcal{L} = g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} (D_i \mathcal{Z}^\dagger D_i \mathcal{Z}) + V(\mathcal{Z}) \right\},$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

- V_0 “hard”; biased towards unitary \mathcal{Z}

- Require gauge and $Z(3)$ invariance

$$\mathcal{Z}(\mathbf{x}) \rightarrow s(\mathbf{x}) \mathcal{Z}(\mathbf{x}) s(\mathbf{x})^\dagger,$$

$$\mathbf{A}(\mathbf{x}) \rightarrow s(\mathbf{x}) (\mathbf{A}(\mathbf{x}) + i\nabla) s(\mathbf{x})^\dagger,$$

$$\mathcal{Z}(\mathbf{x}) \rightarrow e^{2\pi i n/3} \mathcal{Z}(\mathbf{x})$$

and compose Lagrangian as

$$\mathcal{L} = g_3^{-2} \left\{ \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr} (D_i \mathcal{Z}^\dagger D_i \mathcal{Z}) + V(\mathcal{Z}) \right\},$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

- V_0 “hard”; biased towards unitary \mathcal{Z}
- V_1 “soft”; lifts degeneracy and ensures high- T matching to EQCD

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_0(\mathcal{Z}) = c_1 \text{Tr}[\mathcal{Z}^\dagger \mathcal{Z}] + c_2 (\det[\mathcal{Z}] + \det[\mathcal{Z}^\dagger]) \\ + c_3 \text{Tr}[(\mathcal{Z}^\dagger \mathcal{Z})^2]$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_0(\mathcal{Z}) = c_1 \text{Tr}[\mathcal{Z}^\dagger \mathcal{Z}] + c_2 (\det[\mathcal{Z}] + \det[\mathcal{Z}^\dagger]) \\ + c_3 \text{Tr}[(\mathcal{Z}^\dagger \mathcal{Z})^2]$$

- Gives heavy fields their masses

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_0(\mathcal{Z}) = c_1 \text{Tr}[\mathcal{Z}^\dagger \mathcal{Z}] + c_2 (\det[\mathcal{Z}] + \det[\mathcal{Z}^\dagger]) \\ + c_3 \text{Tr}[(\mathcal{Z}^\dagger \mathcal{Z})^2]$$

- Gives heavy fields their masses
- With $c_2 < 0 < c_3$ and $c_2^2 > 9c_1c_3$, V_0 minimized by $Z = \frac{1}{3}v\Omega$ with $\Omega \in \text{SU}(3)$ and

$$v \equiv \frac{3}{4} \left(\frac{-c_2 + \sqrt{c_2^2 - 8c_1c_3}}{c_3} \right)$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_0(\mathcal{Z}) = c_1 \text{Tr}[\mathcal{Z}^\dagger \mathcal{Z}] + c_2 (\det[\mathcal{Z}] + \det[\mathcal{Z}^\dagger]) \\ + c_3 \text{Tr}[(\mathcal{Z}^\dagger \mathcal{Z})^2]$$

- Gives heavy fields their masses
- With $c_2 < 0 < c_3$ and $c_2^2 > 9c_1c_3$, V_0 minimized by $Z = \frac{1}{3}v\Omega$ with $\Omega \in \text{SU}(3)$ and

$$v \equiv \frac{3}{4} \left(\frac{-c_2 + \sqrt{c_2^2 - 8c_1c_3}}{c_3} \right)$$

- Invariant under extra $\text{SU}(3) \times \text{SU}(3)$ symmetry
 $\mathcal{Z}(\mathbf{x}) \rightarrow A\mathcal{Z}(\mathbf{x})B, \quad A, B \in \text{SU}(3)$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_1(\mathcal{Z}) = \tilde{c}_1 \text{Tr} [M^\dagger M] + \tilde{c}_2 (\text{Tr} [M^3] + \text{Tr} [(M^\dagger)^3]) \\ + \tilde{c}_3 \text{Tr} [(M^\dagger M)^2],$$

$$M \equiv \mathcal{Z} - \frac{1}{3} \text{Tr} \mathcal{Z} \mathbf{1} \equiv \mathcal{Z} - \frac{1}{3} L \mathbf{1}$$

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_1(\mathcal{Z}) = \tilde{c}_1 \text{Tr} [M^\dagger M] + \tilde{c}_2 (\text{Tr} [M^3] + \text{Tr} [(M^\dagger)^3]) \\ + \tilde{c}_3 \text{Tr} [(M^\dagger M)^2],$$

$$M \equiv \mathcal{Z} - \frac{1}{3} \text{Tr} \mathcal{Z} \mathbf{1} \equiv \mathcal{Z} - \frac{1}{3} L \mathbf{1}$$

- All $\tilde{c}_i \sim g^0$, but overall $g_3^2 \Rightarrow$ contributions to masses and coupling constants suppressed by g^2 with respect to V_0

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$

$$V_1(\mathcal{Z}) = \tilde{c}_1 \text{Tr}[M^\dagger M] + \tilde{c}_2 (\text{Tr}[M^3] + \text{Tr}[(M^\dagger)^3]) \\ + \tilde{c}_3 \text{Tr}[(M^\dagger M)^2],$$

$$M \equiv \mathcal{Z} - \frac{1}{3} \text{Tr} \mathcal{Z} \mathbf{1} \equiv \mathcal{Z} - \frac{1}{3} L \mathbf{1}$$

- All $\tilde{c}_i \sim g^0$, but overall $g_3^2 \Rightarrow$ contributions to masses and coupling constants suppressed by g^2 with respect to V_0
- Vital for high- T matching to EQCD

$$V(\mathcal{Z}) = V_0(\mathcal{Z}) + g_3^2 V_1(\mathcal{Z})$$


$$V_1(\mathcal{Z}) = \tilde{c}_1 \text{Tr}[M^\dagger M] + \tilde{c}_2 (\text{Tr}[M^3] + \text{Tr}[(M^\dagger)^3]) \\ + \tilde{c}_3 \text{Tr}[(M^\dagger M)^2],$$

$$M \equiv \mathcal{Z} - \frac{1}{3} \text{Tr} \mathcal{Z} \mathbf{1} \equiv \mathcal{Z} - \frac{1}{3} L \mathbf{1}$$

- All $\tilde{c}_i \sim g^0$, but overall $g_3^2 \Rightarrow$ contributions to masses and coupling constants suppressed by g^2 with respect to V_0
- Vital for high- T matching to EQCD
- Assuming $\tilde{c}_3 > 0$ and $\tilde{c}_2^2 < \tilde{c}_1 \tilde{c}_3$, V_1 minimized by $M = 0$, *i.e.* $\mathcal{Z} = \frac{1}{3} L(\mathbf{x}) \mathbf{1}$

- $V(\mathcal{Z})$ minimized by
 - $c_2^2 > 9c_1c_3$: $\mathcal{Z} = \frac{v}{3} e^{2\pi i n/3} \mathbf{1}$
 - $c_2^2 < 9c_1c_3$: $\mathcal{Z} = 0$

- $V(\mathcal{Z})$ minimized by
 - $c_2^2 > 9c_1c_3$: $\mathcal{Z} = \frac{v}{3} e^{2\pi i n/3} \mathbf{1}$
 - $c_2^2 < 9c_1c_3$: $\mathcal{Z} = 0$



Matching to EQCD

- To determine parameters c_i and \tilde{c}_i , consider fluctuations around non-trivial $Z(3)$ minima

$$\mathcal{Z} = e^{2\pi i n/3} \left\{ \frac{1}{3} v \mathbf{1} + g_3 \left[\frac{1}{\sqrt{6}} (\phi + i\chi) \mathbf{1} + (h + ia) \right] \right\}$$

Matching to EQCD

- To determine parameters c_i and \tilde{c}_i , consider fluctuations around non-trivial $Z(3)$ minima

$$\mathcal{Z} = e^{2\pi i n/3} \left\{ \frac{1}{3} v \mathbf{1} + g_3 \left[\frac{1}{\sqrt{6}} (\phi + i\chi) \mathbf{1} + (h + ia) \right] \right\}$$

and write the Lagrangian in terms of the shifted fields

$$\begin{aligned} \mathcal{L} = & V_{\min} + \frac{1}{2} \text{Tr} F_{ij}^2 + \frac{1}{2} [(\partial_i \phi)^2 + m_\phi^2 \phi^2] + \frac{1}{2} [(\partial_i \chi)^2 + m_\chi^2 \chi^2] \\ & + \text{Tr} [(D_i h)^2 + m_h^2 h^2] + \text{Tr} [(D_i a)^2] + V_{\text{int}}(\phi, \chi, h, a) \end{aligned}$$

Matching to EQCD

- To determine parameters c_i and \tilde{c}_i , consider fluctuations around non-trivial $Z(3)$ minima

$$\mathcal{Z} = e^{2\pi i n/3} \left\{ \frac{1}{3} v \mathbf{1} + g_3 \left[\frac{1}{\sqrt{6}} (\phi + i\chi) \mathbf{1} + (h + ia) \right] \right\}$$

and write the Lagrangian in terms of the shifted fields

$$\begin{aligned} \mathcal{L} = & V_{\min} + \frac{1}{2} \text{Tr} F_{ij}^2 + \frac{1}{2} [(\partial_i \phi)^2 + m_\phi^2 \phi^2] + \frac{1}{2} [(\partial_i \chi)^2 + m_\chi^2 \chi^2] \\ & + \text{Tr} [(D_i h)^2 + m_h^2 h^2] + \text{Tr} [(D_i a)^2] + V_{\text{int}}(\phi, \chi, h, a) \end{aligned}$$

- ϕ , χ and h heavy fields, with masses giving c_i 's

$$c_1 = \frac{1}{6}(m_\chi^2 - 3m_\phi^2), \quad c_2 = -m_\chi^2/v,$$

$$c_3 = \frac{3}{4}(m_\chi^2 + 3m_\phi^2)/v^2, \quad m_h^2 = m_\chi^2 + m_\phi^2$$

- At high enough T tunnelings between different vacua exponentially suppressed
 - Integrate out heavy fields ϕ , χ and h to make contact with EQCD

- At high enough T tunnelings between different vacua exponentially suppressed
 - Integrate out heavy fields ϕ , χ and h to make contact with EQCD
- Result: effective theory for a and A_i

$$\mathcal{L}_{\text{light}} = \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr}[(D_i a)^2 + m_a^2 a^2 + \tilde{\lambda} a^4]$$

- At high enough T tunnelings between different vacua exponentially suppressed
 - Integrate out heavy fields ϕ , χ and h to make contact with EQCD
- Result: effective theory for a and A_i

$$\mathcal{L}_{\text{light}} = \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr}[(D_i a)^2 + m_a^2 a^2 + \tilde{\lambda} a^4]$$
- Leading order parameters solely from V_1
 - $\text{SU}(3) \times \text{SU}(3)$ invariance guarantees V_0 does not contribute
 - Result: $\tilde{c}_1 = T + \mathcal{O}(g_3^2)$, $\tilde{c}_3 = \frac{3}{4\pi^2 T} + \mathcal{O}\left(\frac{g_3^2}{T^2}\right)$

- At high enough T tunnelings between different vacua exponentially suppressed
 - Integrate out heavy fields ϕ , χ and h to make contact with EQCD
- Result: effective theory for a and A_i

$$\mathcal{L}_{\text{light}} = \frac{1}{2} \text{Tr} F_{ij}^2 + \text{Tr}[(D_i a)^2 + m_a^2 a^2 + \tilde{\lambda} a^4]$$
- Leading order parameters solely from V_1
 - $\text{SU}(3) \times \text{SU}(3)$ invariance guarantees V_0 does not contribute
 - Result: $\tilde{c}_1 = T + \mathcal{O}(g_3^2)$, $\tilde{c}_3 = \frac{3}{4\pi^2 T} + \mathcal{O}\left(\frac{g_3^2}{T^2}\right)$
- \tilde{c}_2 and v undetermined, but not needed to ensure new theory reproduces *all* EQCD predictions

Z(3) domain walls

- To capture Z(3) physics of the full theory, demand the effective one reproduce leading order domain wall properties

Z(3) domain walls

- To capture Z(3) physics of the full theory, demand the effective one reproduce leading order domain wall properties
- In effective theory, end up minimizing an energy functional expressed in terms of the phases of the eigenvalues of \mathcal{Z}

$$F_{\text{dw}}[\alpha, \beta] \equiv F_{\text{grad}} + F_{\text{soft}} + F_{\text{fluc}} =$$


$$g_3^{-1} (\pi \bar{v} T)^2 \left(\frac{2}{3}\sqrt{T}\right)^3 \int_{-\infty}^{\infty} d\bar{z} \left[(\alpha')^2 + 3(\beta')^2 + U_1 + U_2 \right]$$

$$\text{with } \bar{z} \equiv g_3 \sqrt{T} z, \bar{v} \equiv \frac{v}{T}$$

Z(3) domain walls


- In effective theory, end up minimizing

$$F_{\text{dw}}[\alpha, \beta] \equiv F_{\text{grad}} + F_{\text{soft}} + F_{\text{fluc}} =$$
$$g_3^{-1} (\pi \bar{v} T)^2 \left(\frac{2}{3} \sqrt{T}\right)^3 \int_{-\infty}^{\infty} d\bar{z} \left[(\alpha')^2 + 3(\beta')^2 + U_1 + U_2 \right]$$



- Solve for α, β demanding that domain wall tension and width agree with full theory values
 - $\sigma_{\text{YM}} = \frac{8\pi^2}{9} \frac{T^3}{g(T)}, \Delta z_{\text{YM}} = \frac{\ln(4)-1/2}{g(T)T}$

- Solve for α, β demanding that domain wall tension and width agree with full theory values
 - $\sigma_{\text{YM}} = \frac{8\pi^2}{9} \frac{T^3}{g(T)}, \Delta z_{\text{YM}} = \frac{\ln(4)-1/2}{g(T)T}$
- Result: $v/T = 3.005868, \tilde{c}_2 = 0.118914$



Phase diagram of new theory

- Without perturbative matching, phase diagram parametrized by 6 dimensionless constants
 - With matching, overall scale v known and only g_3^2/v , m_ϕ/m_χ and m_ϕ/v remain

Phase diagram of new theory

- Without perturbative matching, phase diagram parametrized by 6 dimensionless constants
 - With matching, overall scale v known and only g_3^2/v , m_ϕ/m_χ and m_ϕ/v remain
- $V_{\min} = -\frac{v^2}{108g_3^2} (9m_\phi^2 - m_\chi^2) \Rightarrow \text{Z}(3) \text{ invariance}$
spontaneously broken at $m_\phi/m_\chi \lesssim 1/3$

Phase diagram of new theory

- Without perturbative matching, phase diagram parametrized by 6 dimensionless constants
 - With matching, overall scale v known and only g_3^2/v , m_ϕ/m_χ and m_ϕ/v remain
- $V_{\min} = -\frac{v^2}{108g_3^2} (9m_\phi^2 - m_\chi^2) \Rightarrow \text{Z}(3) \text{ invariance}$
spontaneously broken at $m_\phi/m_\chi \lesssim 1/3$
 - At weak coupling, strongly 1st order transition at $m_\phi/m_\chi = 1/3$

Phase diagram of new theory


- Without perturbative matching, phase diagram parametrized by 6 dimensionless constants
 - With matching, overall scale v known and only g_3^2/v , m_ϕ/m_χ and m_ϕ/v remain
- $V_{\min} = -\frac{v^2}{108g_3^2} (9m_\phi^2 - m_\chi^2) \Rightarrow Z(3)$ invariance spontaneously broken at $m_\phi/m_\chi \lesssim 1/3$
 - At weak coupling, strongly 1st order transition at $m_\phi/m_\chi = 1/3$
 - With finite g_3^2/v , expect weakly 1st order fluctuation induced transition

Phase diagram of new theory

- Without perturbative matching, phase diagram parametrized by 6 dimensionless constants
 - With matching, overall scale v known and only g_3^2/v , m_ϕ/m_χ and m_ϕ/v remain
- $V_{\min} = -\frac{v^2}{108g_3^2} (9m_\phi^2 - m_\chi^2) \Rightarrow Z(3)$ invariance spontaneously broken at $m_\phi/m_\chi \lesssim 1/3$
 - At weak coupling, strongly 1st order transition at $m_\phi/m_\chi = 1/3$
 - With finite g_3^2/v , expect weakly 1st order fluctuation induced transition
- In full theory, phase transition known to be weakly 1st order \Rightarrow latter scenario favored

- Numerical simulations needed to study transition and find optimal matching to full theory
 - Match correlation lengths in various channels

- Numerical simulations needed to study transition and find optimal matching to full theory
 - Match correlation lengths in various channels
 - Simulations underway (Kajantie, Kurkela)



Conclusions

- $Z(3)$ invariant effective $3d$ theory constructed for pure $SU(3)$ YM theory
 - Perturbative matching to EQCD ensures correct high temperature predictions
 - Correct domain wall physics built in
 - Phase structure similar to full theory

Conclusions

- Z(3) invariant effective 3d theory constructed for pure SU(3) YM theory
 - Perturbative matching to EQCD ensures correct high temperature predictions
 - Correct domain wall physics built in
 - Phase structure similar to full theory
- Nonperturbative matching to full theory near T_c and nontrivial numerical tests await

Conclusions

- Z(3) invariant effective 3d theory constructed for pure SU(3) YM theory
 - Perturbative matching to EQCD ensures correct high temperature predictions
 - Correct domain wall physics built in
 - Phase structure similar to full theory
- Nonperturbative matching to full theory near T_c and nontrivial numerical tests await
- Possible generalizations: addition of quarks through soft Z(3) breaking terms, higher N_c , ...